

University of Groningen

What makes cyclodextrin glycosyltransferase a transglycosylase

Leemhuis, Reinder Johannes

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2003

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Leemhuis, R. J. (2003). *What makes cyclodextrin glycosyltransferase a transglycosylase*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

REFERENCES

- [1] Stryer,L. (1988) *Biochemistry*, third ed. W. H. Freeman and Company, New York.
- [2] Davies,G.J. & Henrissat,B. (2002) Structural enzymology of carbohydrate-active enzymes: implications for the post-genomic era. *Biochem. Soc. Trans.*, **30**, 291-297.
- [3] Wolfenden,R., Lu,X., & Young,G. (1998) Spontaneous hydrolysis of glycosides. *J. Am. Chem. Soc.*, **120**, 6814-6815.
- [4] Henrissat,B. (1991) A classification of glycosyl hydrolases based on amino acid sequence similarities. *Biochem J.*, **280**, 309-316.
- [5] Carbohydrate-active enzymes server. (1999) (<http://afmb.cnrs-mrs.fr/~cazy/CAZY/index.html>).
- [6] Bourne,Y. & Henrissat,B. (2001) Glycoside hydrolases and glycosyl-transferases: families and functional modules. *Curr. Opin. Struct. Biol.*, **11**, 593-600.
- [7] McCarter,J.D. & Withers,S.G. (1994) Mechanisms of enzymatic glycoside hydrolysis. *Curr. Opin. Struct. Biol.*, **4**, 885-892.
- [8] Shimizu,T., Nakatsu,T., Miyairi,K., Okuno,T., & Kato,H. (2002) Active-site architecture of endopolygalacturonase I from *Stereum purpureum* revealed by crystal structures in native and ligand-bound forms at atomic resolution. *Biochemistry*, **41**, 6651-6659.
- [9] van Santen,Y., Benen,J.A., Schroter,K.H., Kalk,K.H., Armand,S., Visser,J., & Dijkstra,B.W. (1999) 1.68-Å crystal structure of endopolygalacturonase II from *Aspergillus niger* and identification of active site residues by site-directed mutagenesis. *J. Biol. Chem.*, **274**, 30474-30480.

References

- [10] Vocadlo,D.J., Davies,G.J., Laine,R., & Withers,S.G. (2001) Catalysis by hen egg-white lysozyme proceeds via a covalent intermediate. *Nature*, **412**, 835-838.
- [11] Uitdehaag,J.C.M., Mosi,R., Kalk,K.H., van der Veen,B.A., Dijkhuizen,L., Withers,S.G., & Dijkstra,B.W. (1999) X-ray structures along the reaction pathway of cyclodextrin glycosyltransferase elucidate catalysis in the alpha-amylase family. *Nature Struct. Biol.*, **6**, 432-436.
- [12] Terwisscha van Scheltinga,A.C., Armand,S., Kalk,K.H., Isogai,A., Henrissat,B., & Dijkstra,B.W. (1995) Stereochemistry of chitin hydrolysis by a plant chitinase/lysozyme and X-ray structure of a complex with allosamidin: evidence for substrate assisted catalysis. *Biochemistry*, **34**, 15619-15623.
- [13] van Aalten,D.M., Komander,D., Synstad,B., Gaseidnes,S., Peter,M.G., & Eijsink,V.G. (2001) Structural insights into the catalytic mechanism of a family 18 exo-chitinase. *Proc. Natl. Acad. Sci. U.S.A*, **98**, 8979-8984.
- [14] Koshland,D.E. (1953) Stereochemistry and the mechanism of enzymatic reactions. *Biol. Rev. Camb. Philos. Soc.*, **28**, 416-436.
- [15] Wang,Q., Tull,D., Meinke,A., Gilkes,N.R., Warren,R.A., Aebersold,R., & Withers,S.G. (1993) Glu280 is the nucleophile in the active site of *Clostridium thermocellum* CelC, a family A endo-beta-1,4-glucanase. *J. Biol. Chem.*, **268**, 14096-14102.
- [16] Mosi,R., He,S., Uitdehaag,J.C.M., Dijkstra,B.W., & Withers,S.G. (1997) Trapping and characterization of the reaction intermediate in cyclodextrin glycosyltransferase by use of activated substrates and a mutant enzyme. *Biochemistry*, **36**, 9927-9934.
- [17] Davies,G.J., Mackenzie,L., Varrot,A., Dauter,M., Brzozowski,A.M., Schulein,M., & Withers,S.G. (1998) Snapshots along an enzymatic reaction coordinate: analysis of a retaining beta-glycoside hydrolase. *Biochemistry*, **37**, 11707-11713.
- [18] White,A., Tull,D., Johns,K., Withers,S.G., & Rose,D.R. (1996) Crystallographic observation of a covalent catalytic intermediate in a beta-glycosidase. *Nature Struct. Biol.*, **3**, 149-154.

References

- [19] Street,I.P., Kempton,J.B., & Withers,S.G. (1992) Inactivation of a beta-glucosidase through the accumulation of a stable 2-deoxy-2-fluoro-alpha-D-glucopyranosyl-enzyme intermediate: a detailed investigation. *Biochemistry*, **31**, 9970-9978.
- [20] Zechel,D.L. & Withers,S.G. (2000) Glycosidase mechanisms: anatomy of a finely tuned catalyst. *Acc. Chem. Res.* , **33**, 11-18.
- [21] Zechel,D.L. & Withers,S.G. (2001) Dissection of nucleophilic and acid-base catalysis in glycosidases. *Curr. Opin. Chem. Biol.*, **5**, 643-649.
- [22] Rye,C.S. & Withers,S.G. (2000) Glycosidase mechanisms. *Curr. Opin. Chem. Biol.*, **4**, 573-580.
- [23] Janeček,S. (1997) α -Amylase family: molecular biology and evolution. *Prog. Biophys. Molec. Biol.*, **25**, 67-97.
- [24] Svensson,B. (1994) Protein engineering in the alpha-amylase family: catalytic mechanism, substrate specificity, and stability. *Plant Mol. Biol.*, **25**, 141-157.
- [25] Henrissat,B. & Davies,G. (1997) Structural and sequence-based classification of glycoside hydrolases. *Curr. Opin. Struct. Biol.*, **7**, 637-644.
- [26] Kuriki,T. & Imanaka,T. (1999) The concept of the α -amylase family: structural similarity and common catalytic mechanism. *J. Biosci. Bioeng.*, **87**, 557-565.
- [27] Matsuura,Y., Kusunoki,M., Harada,W., & Kakudo,M. (1984) Structure and possible catalytic residues of Taka-amylase A. *J. Biochem. (Tokyo)*, **95**, 697-702.
- [28] Banner,D.W., Bloomer,A.C., Petsko,G.A., Phillips,D.C., Pogson,C.I., Wilson,I.A., Corran,P.H., Furth,A.J., Milman,J.D., Offord,R.E., Priddle,J.D., & Waley,S.G. (1975) Structure of chicken muscle triose phosphate isomerase determined crystallographically at 2.5 angstrom resolution using amino acid sequence data. *Nature*, **255**, 609-614.
- [29] Hocker,B., Jurgens,C., Wilmanns,M., & Sterner,R. (2001) Stability, catalytic versatility and evolution of the (beta-alpha)(8)-barrel fold. *Curr. Opin. Biotechnol.*, **12**, 376-381.

References

- [30] Nakajima,R., Imanaka,T., & Aiba,S. (1986) Comparison of amino acid sequences of eleven different α -amylases. *Appl. Microbiol. Biotechnol.*, **23**, 355-360.
- [31] Uitdehaag,J.C.M., van der Veen,B.A., Dijkhuizen,L., & Dijkstra,B.W. (2002) Catalytic mechanism and product specificity of cyclodextrin glycosyltransferase, a prototypical transglycosylase from the alpha-amylase family. *Enzyme Microb. Technol.*, **30**, 295-304.
- [32] Lawson,C.L., van Montfort,R., Strokopytov,B., Rozeboom,H.J., Kalk,K.H., de Vries,G.E., Penninga,D., Dijkhuizen,L., & Dijkstra,B.W. (1994) Nucleotide sequence and X-ray structure of cyclodextrin glycosyltransferase from *Bacillus circulans* strain 251 in a maltose-dependent crystal form. *J. Mol. Biol.*, **236**, 590-600.
- [33] Penninga,D., van der Veen,B.A., Knegt,R.M., van Hijum,S.A.F.T., Rozeboom,H.J., Kalk,K.H., Dijkstra,B.W., & Dijkhuizen,L. (1996) The raw starch binding domain of cyclodextrin glycosyltransferase from *Bacillus circulans* strain 251. *J. Biol. Chem.*, **271**, 32777-32784.
- [34] Ohdan,K., Kuriki,T., Takata,H., Kaneko,H., & Okada,S. (2000) Introduction of raw starch-binding domains into *Bacillus subtilis* alpha-Amylase by fusion with the starch-binding domain of *Bacillus* cyclomaltodextrin glucanotransferase. *Appl. Environ. Microbiol.*, **66**, 3058-3064.
- [35] Kim,J.S., Cha,S.S., Kim,H.J., Kim,T.J., Ha,N.C., Oh,S.T., Cho,H.S., Cho,M.J., Kim,M.J., Lee,H.S., Kim,J.W., Choi,K.Y., Park,K.H., & Oh,B.H. (1999) Crystal structure of a maltogenic amylase provides insights into a catalytic versatility. *J. Biol. Chem.*, **274**, 26279-26286.
- [36] Park,K.H., Kim,T.J., Cheong,T.K., Kim,J.W., Oh,B.H., & Svensson,B. (2000) Structure, specificity and function of cyclomaltodextrinase, a multispecific enzyme of the alpha-amylase family. *Biochim. Biophys. Acta*, **1478**, 165-185.
- [37] Cheong,K.A., Kim,T.J., Yoon,J.W., Park,C.S., Lee,T.S., Kim,Y.B., Park,K.H., & Kim,J.W. (2002) Catalytic activities of intracellular dimeric neopullulanase on cyclodextrin, acarbose and maltose. *Biotechnol. Appl. Biochem.*, **35**, 27-34.

References

- [38] Kamitori,S., Kondo,S., Okuyama,K., Yokota,T., Shimura,Y., Tonozuka,T., & Sakano,Y. (1999) Crystal structure of *Thermoactinomyces vulgaris* R-47 alpha-amylase II (TVAIL) hydrolyzing cyclodextrins and pullulan at 2.6 Å resolution. *J. Mol. Biol.*, **287**, 907-921.
- [39] Lee,H.S., Kim,M.S., Cho,H.S., Kim,J.I., Kim,T.J., Choi,J.H., Park,C., Lee,H.S., Oh,B.H., & Park,K.H. (2002) Cyclomaltodextrinase, neopullulanase, and maltogenic amylase are nearly indistinguishable from each other. *J. Biol. Chem.* **277**, 21897-21897.
- [40] Strokopytov,B., Knegt,R.M., Penninga,D., Rozeboom,H.J., Kalk,K.H., Dijkhuizen,L., & Dijkstra,B.W. (1996) Structure of cyclodextrin glycosyltransferase complexed with a maltononase inhibitor at 2.6 angstrom resolution. Implications for product specificity. *Biochemistry*, **35**, 4241-4249.
- [41] Uitdehaag,J.C.M., Kalk,K.H., van der Veen,B.A., Dijkhuizen,L., & Dijkstra,B.W. (1999) The cyclization mechanism of cyclodextrin glycosyltransferase (CGTase) as revealed by a gamma-cyclodextrin-CGTase complex at 1.8-Å resolution. *J. Biol. Chem.*, **274**, 34868-34876.
- [42] Uitdehaag,J.C.M., Van Alebeek,G.J., van der Veen,B.A., Dijkhuizen,L., & Dijkstra,B.W. (2000) Structures of maltohexaose and maltoheptaose bound at the donor sites of cyclodextrin glycosyltransferase give insight into the mechanisms of transglycosylation activity and cyclodextrin size specificity. *Biochemistry*, **39**, 7772-7780.
- [43] Strokopytov,B., Penninga,D., Rozeboom,H.J., Kalk,K.H., Dijkhuizen,L., & Dijkstra,B.W. (1995) X-ray structure of cyclodextrin glycosyltransferase complexed with acarbose. Implications for the catalytic mechanism of glycosidases. *Biochemistry*, **34**, 2234-2240.
- [44] Brzozowski,A.M. & Davies,G.J. (1997) Structure of the *Aspergillus oryzae* alpha-amylase complexed with the inhibitor acarbose at 2.0 Å resolution. *Biochemistry*, **36**, 10837-10845.
- [45] Nakamura,A., Haga,K., & Yamane,K. (1993) Three histidine residues in the active center of cyclodextrin glucanotransferase from alkalophilic *Bacillus* sp. 1011: effects of the replacement on pH dependence and transition-state stabilization. *Biochemistry*, **32**, 6624-6631.

References

- [46] Sogaard,M., Kadziola,A., Haser,R., & Svensson,B. (1993) Site-directed mutagenesis of histidine 93, aspartic acid 180, glutamic acid 205, histidine 290, and aspartic acid 291 at the active site and tryptophan 279 at the raw starch binding site in barley alpha-amylase 1. *J. Biol. Chem.*, **268**, 22480-22484.
- [47] Ly,H.D. & Withers,S.G. (1999) Mutagenesis of glycosidases. *Annu. Rev. Biochem.*, **68**, 487-522.
- [48] Libessart,N. & Preiss,J. (1998) Arginine residue 384 at the catalytic center is important for branching enzyme II from maize endosperm. *Arch. Biochem. Biophys.*, **360**, 135-141.
- [49] Mirza,O., Skov,L.K., Remaud-Simeon,M., Potocki,d.M., Albenne,C., Monsan,P., & Gajhede,M. (2001) Crystal structures of amylosucrase from *Neisseria polysaccharea* in complex with D-glucose and the active site mutant Glu328Gln in complex with the natural substrate sucrose. *Biochemistry*, **40**, 9032-9039.
- [50] Watanabe,K., Hata,Y., Kizaki,H., Katsube,Y., & Suzuki,Y. (1997) The refined crystal structure of *Bacillus cereus* oligo-1,6-glucosidase at 2.0 Å resolution: structural characterization of proline-substitution sites for protein thermostabilization. *J. Mol. Biol.*, **269**, 142-153.
- [51] Katsuya,Y., Mezaki,Y., Kubota,M., & Matsuura,Y. (1998) Three-dimensional structure of *Pseudomonas* isoamylase at 2.2 Å resolution. *J. Mol. Biol.*, **281**, 885-897.
- [52] Abad,M.C., Binderup,K., Rios-Steiner,J., Arni,R.K., Preiss,J., & Geiger,J.H. (2002) The X-ray crystallographic structure of *Escherichia coli* branching enzyme. *J. Biol. Chem.*, **277**, 42164-42170.
- [53] Leemhuis,H., Dijkstra,B.W., & Dijkhuizen,L. (2002) Mutations converting cyclodextrin glycosyltransferase from a transglycosylase into a starch hydrolase. *FEBS Lett.*, **514**, 189-192.
- [54] Kuriki,T., Kaneko,H., Yanase,M., Takata,H., Shimada,J., Handa,S., Takada,T., Umeyama,H., & Okada,S. (1996) Controlling substrate preference and transglycosylation activity of neopullulanase by manipulating steric constraint and hydrophobicity in active center. *J. Biol. Chem.*, **271**, 17321-17329.

References

- [55] Roujeinikova,A., Raasch,C., Burke,J., Baker,P.J., Liebl,W., & Rice,D.W. (2001) The crystal structure of *Thermotoga maritima* maltosyltransferase and its implications for the molecular basis of the novel transfer specificity. *J. Mol. Biol.*, **312**, 119-131.
- [56] Dauter,Z., Dauter,M., Brzozowski,A.M., Christensen,S., Borchert,T.V., Beier,L., Wilson,K.S., & Davies,G.J. (1999) X-ray structure of Novamyl, the five-domain "maltogenic" alpha-amylase from *Bacillus stearothermophilus*: maltose and acarbose complexes at 1.7Å resolution. *Biochemistry*, **38**, 8385-8392.
- [57] Christophersen,C., Otzen,D.E., Norman,B.E., Christensen,S., & Schäfer,T. (1998) Enzymatic characterization of Novamyl, a thermostable α -amylase. *Starch/Stärke*, **50**, 39-45.
- [58] Wind,R.D., Uitdehaag,J.C.M., Buitelaar,R.M., Dijkstra,B.W., & Dijkhuizen,L. (1998) Engineering of cyclodextrin product specificity and pH optima of the thermostable cyclodextrin glycosyltransferase from *Thermoanaerobacterium thermosulfurigenes* EM1. *J. Biol. Chem.*, **273**, 5771-5779.
- [59] Hehre,E.J., Genghof,D.S., & Okada,G. (1971) The alpha-amylases as glycosylases, with wider catalytic capacities than envisioned or explained by their representation as hydrolases. *Arch. Biochem. Biophys.*, **142**, 382-393.
- [60] Brayer,G.D., Sidhu,G., Maurus,R., Rydberg,E.H., Braun,C., Wang,Y., Nguyen,N.T., Overall,C.M., & Withers,S.G. (2000) Subsite mapping of the human pancreatic alpha-amylase active site through structural, kinetic, and mutagenesis techniques. *Biochemistry*, **39**, 4778-4791.
- [61] Aghajari,N., Roth,M., & Haser,R. (2002) Crystallographic evidence of a transglycosylation reaction: ternary complexes of a psychrophilic alpha-amylase. *Biochemistry*, **41**, 4273-4280.
- [62] Saab-Rincon,G., del Rio,G., Santamaria,R.I., Lopez-Munguia,A., & Soberon,X. (1999) Introducing transglycosylation activity in a liquefying alpha-amylase. *FEBS Lett.*, **453**, 100-106.

References

- [63] Matsui,I., Ishikawa,K., Miyairi,S., Fukui,S., & Honda,K. (1991) An increase in the transglycosylation activity of *Saccharomycopsis* alpha-amylase altered by site-directed mutagenesis. *Biochim. Biophys. Acta*, **1077**, 416-419.
- [64] Matsui,I., Yoneda,S., Ishikawa,K., Miyairi,S., Fukui,S., Umeyama,H., & Honda,K. (1994) Roles of the aromatic residues conserved in the active center of *Saccharomycopsis* alpha-amylase for transglycosylation and hydrolysis activity. *Biochemistry* , **33**, 451-458.
- [65] Qian,M., Haser,R., Buisson,G., Duee,E., & Payan,F. (1994) The active center of a mammalian alpha-amylase. Structure of the complex of a pancreatic alpha-amylase with a carbohydrate inhibitor refined to 2.2-Å resolution. *Biochemistry*, **33**, 6284-6294.
- [66] French,D., Pulley,A.O., Effenberger,J.A., Rougvie,M.A., & Abdullah, M. (1965) Studies on the Schardinger dextrans. XII. The molecular size and structure of the delta-, epsilon-, zeta-, and eta-dextrans. *Arch. Biochem. Biophys.*, **111**, 153-160.
- [67] Terada,Y., Yanase,M., Takata,H., Takaha,T., & Okada,S. (1997) Cyclodextrins are not the major cyclic alpha-1,4-glucans produced by the initial action of cyclodextrin glucanotransferase on amylose. *J. Biol. Chem.*, **272**, 15729-15733.
- [68] Penninga,D., Strokopytov,B., Rozeboom,H.J., Lawson,C.L., Dijkstra, B.W., Bergsma,J., & Dijkhuizen,L. (1995) Site-directed mutations in tyrosine 195 of cyclodextrin glycosyltransferase from *Bacillus circulans* strain 251 affect activity and product specificity. *Biochemistry*, **34**, 3368-3376.
- [69] Klein,C. & Schulz,G.E. (1991) Structure of cyclodextrin glycosyltransferase refined at 2.0 Å resolution. *J. Mol. Biol.*, **217**, 737-750.
- [70] Knegt,R.M., Wind,R.D., Rozeboom,H.J., Kalk,K.H., Buitelaar,R.M., Dijkhuizen,L., & Dijkstra,B.W. (1996) Crystal structure at 2.3 Å resolution and revised nucleotide sequence of the thermostable cyclodextrin glycosyltransferase from *Thermonaerobacterium thermosulfurigenes* EM1. *J. Mol. Biol.*, **256**, 611-622.

References

- [71] Kubota,M., Matsuura,Y., Sakai,S., & Katsube,Y. (1991) Molecular structure of *B. stearothermophilus* cyclodextrin glucanotransferase and analysis of substrate binding site. *Denpun Kagaku*, **38**, 141-146.
- [72] Harata,K., Haga,K., Nakamura,A., Aoyagi,M., & Yamane,K. (1996) X-ray structure of cyclodextrin glucanotransferase from alkalophilic *Bacillus* sp. 1011. Comparision of two independent molecules at 1.8 Å resolution. *Acta Crystallogr.*, **D52**, 1136-1145.
- [73] Schmidt,A.K., Cottaz,S., Driguez,H., & Schulz,G.E. (1998) Structure of cyclodextrin glycosyltransferase complexed with a derivative of its main product beta-cyclodextrin. *Biochemistry*, **37**, 5909-5915.
- [74] Leemhuis,H., Uitdehaag,J.C.M., Rozeboom,H.J., Dijkstra,B.W., & Dijkhuizen,L. (2002) The remote substrate binding subsite -6 in cyclodextrin glycosyltransferase controls the transferase activity of the enzyme via an induced-fit mechanism. *J. Biol. Chem.*, **277**, 1113-1119.
- [75] van der Veen,B.A., Uitdehaag,J.C.M., Penninga,D., Van Alebeek,G.J., Smith,L.M., Dijkstra,B.W., & Dijkhuizen,L. (2000) Rational design of cyclodextrin glycosyltransferase from *Bacillus circulans* strain 251 to increase alpha-cyclodextrin production. *J. Mol. Biol.*, **296**, 1027-1038.
- [76] van der Veen,B.A., Uitdehaag,J.C.M., Dijkstra,B.W., & Dijkhuizen,L. (2000) The role of arginine 47 in the cyclization and coupling reactions of cyclodextrin glycosyltransferase from *Bacillus circulans* strain 251 Implications for product inhibition and product specificity. *Eur. J. Biochem.*, **267**, 3432-3441.
- [77] Parsiegla,G., Schmidt,A.K., & Schulz,G.E. (1998) Substrate binding to a cyclodextrin glycosyltransferase and mutations increasing the gamma-cyclodextrin production. *Eur. J. Biochem.*, **255**, 710-717.
- [78] Beier,L., Svendsen,A., Andersen,C., Frandsen,T.P., Borchert,T.V., & Cherry,J.R. (2000) Conversion of the maltogenic alpha-amylase Novamyl into a CGTase. *Protein Eng.*, **13**, 509-513.
- [79] Roujeinikova,A., Raasch,C., Sedelnikova,S., Liebl,W., & Rice,D.W. (2002) Crystal Structure of *Thermotoga maritima* 4-alpha-Glucanotransferase and its Acarbose Complex: Implications for Substrate Specificity and Catalysis. *J. Mol. Biol.*, **321**, 149-162.

References

- [80] Kuriki,T., Takata,H., Okada,S., & Imanaka,T. (1991) Analysis of the active center of *Bacillus stearothermophilus* neopullulanase. *J. Bacteriol.*, **173**, 6147-6152.
- [81] Takata,H., Kuriki,T., Okada,S., Takesada,Y., Iizuka,M., Minamiura,N., & Imanaka,T. (1992) Action of neopullulanase. Neopullulanase catalyzes both hydrolysis and transglycosylation at alpha-(1,4)- and alpha-(1,6)-glucosidic linkages. *J. Biol. Chem.*, **267**, 18447-18452.
- [82] Hashimoto,Y., Yamamoto,T., Fujiwara,S., Takagi,M., & Imanaka,T. (2001) Extracellular synthesis, specific recognition, and intracellular degradation of cyclomaltodextrins by the hyperthermophilic archaeon *Thermococcus* sp. strain B1001. *J. Bacteriol.*, **183**, 5050-5057.
- [83] Fiedler,G., Pajatsch,M., & Bock,A. (1996) Genetics of a novel starch utilisation pathway present in *Klebsiella oxytoca*. *J. Mol. Biol.*, **256**, 279-291.
- [84] Pajatsch,M., Andersen,C., Mathes,A., Bock,A., Benz,R., & Engelhardt,H. (1999) Properties of a cyclodextrin-specific, unusual porin from *Klebsiella oxytoca*. *J. Biol. Chem.*, **274**, 25159-25166.
- [85] Pajatsch,M., Gerhart,M., Peist,R., Horlacher,R., Boos,W., & Bock,A. (1998) The periplasmic cyclodextrin binding protein CymE from *Klebsiella oxytoca* and its role in maltodextrin and cyclodextrin transport. *J. Bacteriol.*, **180**, 2630-2635.
- [86] van der Veen,B.A., Uitdehaag,J.C.M., Dijkstra,B.W., & Dijkhuizen,L. (2000) Engineering of cyclodextrin glycosyltransferase reaction and product specificity. *Biochim. Biophys. Acta*, **1543**, 336-360.
- [87] Cha,H.J., Yoon,H.G., Kim,Y.W., Lee,H.S., Kim,J.W., Kweon,K.S., Oh,B.H., & Park,K.H. (1998) Molecular and enzymatic characterization of a maltogenic amylase that hydrolyzes and transglycosylates acarbose. *Eur. J. Biochem.*, **253**, 251-262.
- [88] Kim,T.J., Park,C.S., Cho,H.Y., Cha,S.S., Kim,J.S., Lee,S.B., Moon,T.W., Kim,J.W., Oh,B.H., & Park,K.H. (2000) Role of the glutamate 332 residue in the transglycosylation activity of *thermus* maltogenic amylase. *Biochemistry*, **39**, 6773-6780.

References

- [89] Nakayama,A., Yamamoto,K., & Tabata,S. (2001) Identification of the catalytic residues of bifunctional glycogen debranching enzyme. *J. Biol. Chem.*, **276**, 28824-28828.
- [90] Hatada,Y., Igarashi,K., Ozaki,K., Ara,K., Hitomi,J., Kobayashi,T., Kawai,S., Watabe,T., & Ito,S. (1996) Amino acid sequence and molecular structure of an alkaline amylopullulanase from *Bacillus* that hydrolyzes alpha-1,4 and alpha-1,6 linkages in polysaccharides at different active sites. *J. Biol. Chem.*, **271**, 24075-24083.
- [91] Lim,W.J., Park,S.R., Cho,S.J., Kim,M.K., Ryu,S.K., Hong,S.Y., Seo,W.T., Kim,H., & Yun,H.D. (2001) Cloning and characterization of an intracellular isoamylase gene from *Pectobacterium chrysanthemi* PY35. *Biochem. Biophys. Res. Commun.*, **287**, 348-354.
- [92] Duffner,F., Bertoldo,C., Andersen,J.T., Wagner,K., & Antranikian,G. (2000) A new thermoactive pullulanase from *Desulfurococcus mucosus*: cloning, sequencing, purification, and characterization of the recombinant enzyme after expression in *Bacillus subtilis*. *J. Bacteriol.*, **182**, 6331-6338.
- [93] Preiss,J. (1984) Bacterial glycogen synthesis and its regulation. *Annu. Rev. Microbiol.*, **38**, 419-458.
- [94] Nakada,T., Maruta,K., Tsusaki,K., Kubota,M., Chaen,H., Sugimoto,T., Kurimoto,M., & Tsujisaka,Y. (1995) Purification and properties of a novel enzyme, maltooligosyl trehalose synthase, from *Arthrobacter* sp. Q36. *Biosci. Biotechnol. Biochem.*, **59**, 2210-2214.
- [95] Nakada,T., Maruta,K., Mitsuzumi,H., Kubota,M., Chaen,H., Sugimoto,T., Kurimoto,M., & Tsujisaka,Y. (1995) Purification and characterization of a novel enzyme, maltooligosyl trehalose trehalohydrolase, from *Arthrobacter* sp. Q36. *Biosci. Biotechnol. Biochem.*, **59**, 2215-2218.
- [96] Gueguen,Y., Rolland,J.L., Schroeck,S., Flament,D., Defretin,S., Saniez,M.H., & Dietrich,J. (2001) Characterization of the maltooligosyl trehalose synthase from the thermophilic archaeon *Sulfolobus acidocaldarius*. *FEMS Microbiol. Lett.*, **194**, 201-206.

References

- [97] Buttcher,V., Welsh,T., Willmitzer,L., & Kossmann,J. (1997) Cloning and characterization of the gene for amylosucrase from *Neisseria polysaccharea*: production of a linear alpha-1,4-glucan. *J. Bacteriol.*, **179**, 3324-3330.
- [98] Potocki,d.M., Remaud-Simeon,M., Willemot,R.M., Sarcabal,P., Planchot,V., & Monsan,P. (2000) Amylosucrase from *Neisseria polysaccharea*: novel catalytic properties. *FEBS Lett.*, **471**, 219-223.
- [99] Skov,L.K., Mirza,O., Henriksen,A., De Montalk,G.P., Remaud-Simeon,M., Sarcabal,P., Willemot,R.M., Monsan,P., & Gajhede,M. (2001) Amylosucrase, a glucan-synthesizing enzyme from the alpha-amylase family. *J. Biol. Chem.*, **276**, 25273-25278.
- [100] Penninga,D. (1996) Protein engineering of cyclodextrin glycosyl-transferase from *Bacillus circulans* strain 251. Ph. D. thesis, University of Groningen, the Netherlands.
- [101] Wind,R.D. (1997) Starch-converting enzymes from thermophilic microorganisms. Ph. D. thesis, University of Groningen, the Netherlands.
- [102] van der Veen,B.A. (2000) Engineering reaction and product specificity of cyclodextrin glycosyltransferase from *Bacillus circulans* strain 251. Ph. D. thesis, University of Groningen, the Netherlands.
- [103] Uitdehaag,J.C.M. (2000) How cyclodextrin glycosyltransferase works: the atomic basis of catalysis in an enzyme of the α -amylase family. Ph. D. thesis, University of Groningen, the Netherlands.
- [104] Davies,G.J., Wilson,K.S., & Henrissat,B. (1997) Nomenclature for sugar-binding subsites in glycosyl hydrolases. *Biochem. J.*, **321**, 557-559.
- [105] van der Maarel,M.J.E.C., van der Veen,B.A., Uitdehaag,J.C.M., Leemhuis,H. & Dijkhuizen,L. (2002) Properties and applications of starch-converting enzymes of the α -amylase family. *J. Biotechnol.*, **94**, 137-155.
- [106] Lee,K.C.P. & Tao,B.Y. (1995) A kinetic study of cyclodextrin glycosyltransferase: substrate and product inhibition. *Biotechnol. Appl. Biochem.*, **21**, 111-121.

References

- [107] Pedersen,S., Dijkhuizen,L., Dijkstra,B.W., Jensen,B.F., & Jørgensen, S.T. (1995) A better enzyme for cyclodextrins. *Chemtech*, **12**, 19-25.
- [108] Ferey-Roux,G., Perrier,J., Forest,E., Marchis-Mouren,G., Puigserver,A., & Santimone,M. (1998) The human pancreatic alpha-amylase isoforms: isolation, structural studies and kinetics of inhibition by acarbose. *Biochim. Biophys. Acta*, **1388**, 10-20.
- [109] Koukietolo,R., Berre-Anton,V., Desseaux,V., Moreau,Y., Rouge,P., Marchis-Mouren,G., & Santimone,M. (1999) Mechanism of porcine pancreatic alpha-amylase inhibition of amylose and maltopentaose hydrolysis by kidney bean (*Phaseolus vulgaris*) inhibitor and comparison with that by acarbose. *Eur. J. Biochem.*, **265**, 20-26.
- [110] Alkazaz,M., Desseaux,V., Marchis-Mouren,G., Payan,F., Forest,E., & Santimone,M. (1996) The mechanism of porcine pancreatic alpha-amylase. Kinetic evidence for two additional carbohydrate-binding sites. *Eur. J. Biochem.*, **241**, 787-796.
- [111] Al Kazaz,M., Desseaux,V., Marchis-Mouren,G., Prodanov,E., & Santimone,M. (1998) The mechanism of porcine pancreatic alpha-amylase. Inhibition of maltopentaose hydrolysis by acarbose, maltose and maltotriose. *Eur. J. Biochem.*, **252**, 100-107.
- [112] Koukietolo,R., Desseaux,V., Moreau,Y., Marchis-Mouren,G., & Santimone,M. (2001) Mechanism of porcine pancreatic alpha-amylase. Inhibition of amylose and maltopentaose hydrolysis by alpha-, beta- and gamma-cyclodextrins. *Eur. J. Biochem.*, **268**, 841-848.
- [113] Kim,M.J., Lee,S.B., Lee,H.S., Lee,S.Y., Baek,J.S., Kim,D., Moon,T.W., Robyt,J.F., & Park,K.H. (1999) Comparative study of the inhibition of alpha-glucosidase, alpha-amylase, and cyclomaltodextrin glucanotransferase by acarbose, isoacarboses, and acarviosine-glucose. *Arch. Biochem. Biophys.*, **371**, 277-283.
- [114] Mosi,R., Sham,H., Uitdehaag,J.C.M., Ruiterkamp,R., Dijkstra,B.W., & Withers,S.G. (1998) Reassessment of acarbose as a transition state analogue inhibitor of cyclodextrin glycosyltransferase. *Biochemistry*, **37**, 17192-17198.

References

- [115] Haeckel, K. & Bahl, H. (1989) Cloning and expression of the thermostable α -amylase gene from *Clostridium thermosulfurigenes* (DSM 3896) in *Escherichia coli*. *FEMS Microbiol. Lett.*, **60**, 333-338.
- [116] Smith, H., de Jong, A., Bron, S., & Venema, G. (1988) Characterization of signal-sequence-coding regions selected from the *Bacillus subtilis* chromosome. *Gene*, **70**, 351-361.
- [117] Bron, S. (1990) Plasmids. In *Modern microbiological methods for Bacillus* (Harwood, C.R. & Cutting, S.M., Eds), pp. 146-147. John Wiley & Sons, New York/Chichester.
- [118] Vikmon, M. (1982) Rapid and simple spectrophotometric method for determination of microamounts of cyclodextrins (Szejtli, J. Ed.), pp 64-74. Proceedings of the first international symposium on cyclodextrins. Reidel Publishing Co. Dordrecht, The Netherlands.
- [119] Cornish-Bowden, A. (1995) *Fundamentals of enzyme kinetics*. Portland Press, London.
- [120] Wind, R.D., Liebl, W., Buitelaar, R.M., Penninga, D., Spreinat, A., Dijkhuizen, L., & Bahl, H. (1995) Cyclodextrin formation by the thermostable α -amylase of *Thermoanaerobacterium thermosulfurigenes* EM1 and reclassification of the enzyme as a cyclodextrin glycosyltransferase. *Appl. Environ. Microbiol.*, **61**, 1257-1265.
- [121] Nakamura, A., Haga, K., & Yamane, K. (1994) The transglycosylation reaction of cyclodextrin glucanotransferase is operated by a Ping-Pong mechanism. *FEBS Lett.*, **337**, 66-70.
- [122] van der Veen, B.A., Van Alebeek, G.J., Uitdehaag, J.C.M., Dijkstra, B.W., & Dijkhuizen, L. (2000) The three transglycosylation reactions catalyzed by cyclodextrin glycosyltransferase from *Bacillus circulans* (strain 251) proceed via different kinetic mechanisms. *Eur. J. Biochem.*, **267**, 658-665.
- [123] Nakamura, A., Haga, K., & Yamane, K. (1994) Four aromatic residues in the active center of cyclodextrin glucanotransferase from alkalophilic *Bacillus* sp. 1011: effects of replacements on substrate binding and cyclization characteristics. *Biochemistry*, **33**, 9929-9936.
- [124] van der Veen, B.A., Leemhuis, H., Kralj, S., Uitdehaag, J.C.M., Dijkstra, B.W., & Dijkhuizen, L. (2001) Hydrophobic amino acid

References

- residues in the acceptor binding site are main determinants for reaction mechanism and specificity of cyclodextrin glycosyltransferase. *J. Biol. Chem.*, **276**, 44557-44562.
- [125] Bateman,A., Birney,E., Cerruti,L., Durbin,R., Eddy,S.R., Griffiths-Jones,S., Howe,K.L., Marshall,M., & Sonnhammer,E.L. (2002) The Pfam protein families database. *Nucleic Acids Res.*, **30**, 276-280.
- [126] Vihinen,M., Ollikka,P., Niskanen,J., Meyer,P., Suominen,I., Karp,M., Holm,L., Knowles,J., & Mantsala,P. (1990) Site-directed mutagenesis of a thermostable alpha-amylase from *Bacillus stearothermophilus*: putative role of three conserved residues. *J. Biochem. (Tokyo)*, **107**, 267-272.
- [127] Otwinowski,Z. (1993) In *Data collection and Processing* (Sawyer,L., Isaacs,N., & Bailey,S., Eds), pp. 56-62. SERC Laboratory, Daresbury, UK.
- [128] Brünger,A.T., Adams,P.D., Clore,G.M., DeLano,W.L., Gros,P., Grosse-Kunstleve,R.W., Jiamg,J.-S., Kuszewski,J., Nilges,M., Pannu,N.S., Read,R.J., Rice,L.M., Simonson,T., & Warren,G.L. (1998) Crystallography & NMR system: A new software suite for macromolecular structure determination. *Acta Cryst.*, **D54**, 905-921.
- [129] Jones,T.A., Zou,J.Y., Cowan,S.W., & Kjeldgaard,M. (1991) Improved methods for building protein models in electron density maps and the location of errors in these models. *Acta Crystallogr.*, **A47**, 110-119.
- [130] Meissner,P.S., Sisk,W.P., & Berman,M.L. (1987) Bacteriophage lambda cloning system for the construction of directional cDNA libraries. *Proc. Natl. Acad. Sci. U.S.A.*, **84**, 4171-4175.
- [131] Leemhuis,H., Dijkstra,B.W., & Dijkhuizen,L. (2003) *Thermoanaerobacterium thermosulfurigenes* cyclodextrin glycosyltransferase: mechanism and kinetics of inhibition by acarbose and cyclodextrins. *Eur. J. Biochem.*, **270**, 155-162.
- [132] Sambrook,J., Frisch,E.J., & Maniatis,T. (1989) *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, New York.

References

- [133] Guex,N. & Peitsch,M.C. (1997) SWISS-MODEL and the Swiss-PdbViewer: an environment for comparative protein modeling. *Electrophoresis*, **18**, 2714-2723.
- [134] Nakayama,A., Yamamoto,K., & Tabata,S. (2001) The role of Asp-224 in conserved sequence region I of yeast glycogen debranching enzyme. *Applied Biological Science*, **7**, 37-47.
- [135] Knegtel,R.M., Strokopytov,B., Penninga,D., Faber,O.G., Rozeboom,H.J., Kalk,K.H., Dijkhuizen,L., & Dijkstra,B.W. (1995) Crystallographic studies of the interaction of cyclodextrin glycosyltransferase from *Bacillus circulans* strain 251 with natural substrates and products. *J. Biol. Chem.*, **270**, 29256-29264.
- [136] Kim,Y.H., Bae,K.H., Kim,T.J., Park,K.H., Lee,H.S., & Byun,S.M. (1997) Effect on product specificity of cyclodextrin glycosyltransferase by site-directed mutagenesis. *Biochem. Mol. Biol. Int.*, **41**, 227-234.
- [137] Uitdehaag,J.C.M., van der Veen,B.A., Dijkhuizen,L., Elber,R., & Dijkstra,B.W. (2001) Enzymatic circularization of a malto-octaose linear chain studied by stochastic reaction path calculations on cyclodextrin glycosyltransferase. *Proteins:Struct. Funct. Genet.*, **43**, 327-335.
- [138] Kabsch,W. (1988) Evaluation of single-crystal diffraction data from a positive-sensitive detector. *J. Appl. Cryst.*, **21**, 916-924.
- [139] Murray,V. (1989) Improved double-stranded DNA sequencing using the linear polymerase chain reaction. *Nucleic Acids Res.*, **17**, 8889.
- [140] Bender,H. (1990) Studies of the mechanism of the cyclisation reaction catalysed by the wildtype and a truncated alpha-cyclodextrin glycosyltransferase from *Klebsiella pneumoniae* strain M 5 al, and the beta-cyclodextrin glycosyltransferase from *Bacillus circulans* strain 8. *Carbohydr. Res.*, **206**, 257-267.
- [141] Hanahan,D. (1983) Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.*, **166**, 557-580.

References

- [142] van der Vossen,J.M., Kodde,J., Haandrikman,A.J., Venema,G., & Kok,J. (1992) Characterization of transcription initiation and termination signals of the proteinase genes of *Lactococcus lactis* Wg2 and enhancement of proteolysis in *L. lactis*. *Appl. Environ. Microbiol.*, **58**, 3142-3149.
- [143] Voss,H., Wirkner,U., Zimmermann,J., Erfle,H., Hewitt,N., Rupp,T., Stegemann,J., & Ansorge,W. (1992) New procedure for automated DNA sequencing with multiple internal labeling by fluorescent dUTP. *Methods Mol. Cell Biol.*, **3**, 30-34.
- [144] Zimmermann,J., Voss,H., Schwager,C., Stegemann,J., Erfle,H., Stucky,K., Kristensen,T., & Ansorge,W. (1990) A simplified protocol for fast plasmid DNA sequencing. *Nucleic Acids Res.*, **18**, 1067.
- [145] Sundberg,L. & Porath,J. (1974) Preparation of adsorbents for biospecific affinity chromatography. *J. Chromatogr.*, **90**, 87-98.
- [146] Bernfeld,P. (1955) Amylases, α and β . *Meth. Enzymol.*, **1**, 149-158.
- [147] Norman,B.E. & Jørgensen S.T. (1992) *Thermoanaerobacter* sp. CGTase: its properties and application. *Denpun Kagaku*, **39**, 101-108.
- [148] Klein,C., Hollender,J., Bender,H., & Schulz,G.E. (1992) Catalytic center of cyclodextrin glycosyltransferase derived from X-ray structure analysis combined with site-directed mutagenesis. *Biochemistry*, **31**, 8740-8746.
- [149] Fujiwara,S., Kakihara,H., Sakaguchi,K., & Imanaka,T. (1992) Analysis of mutations in cyclodextrin glucanotransferase from *Bacillus stearothermophilus* which affect cyclization characteristics and thermostability. *J. Bacteriol.*, **174**, 7478-7481.
- [150] Sin,K.A., Nakamura,A., Masaki,H., Matsuura,Y., & Uozumi,T. (1994) Replacement of an amino acid residue of cyclodextrin glucanotransferase of *Bacillus ohbensis* doubles the production of gamma-cyclodextrin. *J. Biotechnol.*, **32**, 283-288.
- [151] Leemhuis,H., Dijkstra,B.W., & Dijkhuizen,L. *In preparation*.

References

- [152] Alcalde,M., Plou,F.J., Andersen,C., Martin,M.T., Pedersen,S., & Ballesteros,A. (1999) Chemical modification of lysine side chains of cyclodextrin glycosyltransferase from *Thermoanaerobacter* causes a shift from cyclodextrin glycosyltransferase to alpha-amylase specificity. *FEBS Lett*, **445**, 333-337.
- [153] Somogyi,M. (1952) Notes on sugar determination. *J. Biol. Chem.*, **195**, 19-23.
- [154] Nelson,N. (2002) A photometric adaption on the Somogyi method for the determination of glucose. *J. Biol. Chem.*, **153**, 375-380.
- [155] Berman,H.M., Westbrook,J., Feng,Z., Gilliland,G., Bhat,T.N., Weissig,H., Shindyalov,I.N., & Bourne,P.E. (2000) The Protein Data Bank. *Nucleic Acids Res.*, **28**, 235-242.
- [156] Leung,D.W., Chen,E., & Goeddel,D.V. (1989) A method for random mutagenesis of a defined DNA segment using a modified polymerase chain reaction. *Technique*, **1**, 11-15.
- [157] Mattsson,P., Battchikova,N., Sippola,K., & Korpela,T. (1995) The role of histidine residues in the catalytic act of cyclomaltodextrin glucanotransferase from *Bacillus circulans* var. alkalophilus. *Biochim. Biophys. Acta*, **1247**, 97-103.
- [158] Nielsen,J.E., Beier,L., Otzen,D., Borchert,T.V., Frantzen,H.B., Andersen,K.V., & Svendsen,A. (1999) Electrostatics in the active site of an alpha-amylase. *Eur. J. Biochem.*, **264**, 816-824.
- [159] Crabb,W.D. & Mitchinson,C. (1997) Enzymes involved in the processing of starch to sugars. *Tibtech*, **15**, 349-352.
- [160] Diderichsen,B. & Christiansen,L. (1988) Cloning of a maltogenic alpha-amylase from *Bacillus stearothermophilus*. *FEMS Microbiol. Lett.*, **56**, 53-60.
- [161] Lazzaroni,J.C. & Portalier,R. (1979) Isolation and preliminary characterization of periplasmic-leaky mutants of *Escherichia coli* K-12. *FEMS Microbiol. Lett.*, **5**, 411-416.

References

- [162] Uitdehaag,J.C.M. & Dijkstra,B.W. (1998) A strategy for engineering thermostability: the case of cyclodextrin glycosyltransferase. In *Stability and stabilization of biocatalysts* (Ballesteros,A., ed), pp. 317-323. Elsevier.
- [163] Rashid,N., Cornista,J., Ezaki,S., Fukui,T., Atomi,H., & Imanaka,T. (2002) Characterization of an archaeal cyclodextrin glucanotransferase with a novel C-terminal domain. *J. Bacteriol.*, **184**, 777-784.
- [164] Yamamoto,K., Shiraki,K., Fujiwara,S., Takagi,M., Fukui,K., & Imanaka,T. (1999) *In vitro* heat effect on functional and conformational changes of cyclodextrin glucanotransferase from hyperthermophilic archaea. *Biochem. Biophys. Res. Commun.*, **265**, 57-61.
- [165] Tachibana,Y., Kuramura,A., Shirasaka,N., Suzuki,Y., Yamamoto,T., Fujiwara,S., Takagi,M., & Imanaka,T. (1999) Purification and characterization of an extremely thermostable cyclomaltodextrin glucanotransferase from a newly isolated hyperthermophilic archaeon, a *Thermococcus* sp. *Appl. Environ. Microbiol.*, **65**, 1991-1997.
- [166] Savchenko,A., Vieille,C., Kang,S., & Zeikus,J.G. (2002) *Pyrococcus furiosus* alpha-amylase is stabilized by calcium and zinc. *Biochemistry*, **41**, 6193-6201.
- [167] Igarashi,K., Hatada,Y., Ikawa,K., Araki,H., Ozawa,T., Kobayashi,T., Ozaki,K., & Ito,S. (1998) Improved thermostability of a *Bacillus* alpha-amylase by deletion of an arginine-glycine residue is caused by enhanced calcium binding. *Biochem. Biophys. Res. Commun.*, **248**, 372-377.
- [168] Declerck,N., Joyet,P., Gaillardin,C., & Masson,J.M. (1990) Use of amber suppressors to investigate the thermostability of *Bacillus licheniformis* alpha-amylase. Amino acid replacements at 6 histidine residues reveal a critical position at His-133. *J. Biol. Chem.*, **265**, 15481-15488.
- [169] Declerck,N., Machius,M., Chambert,R., Wiegand,G., Huber,R., & Gaillardin,C. (1997) Hyperthermostable mutants of *Bacillus licheniformis* alpha-amylase: thermodynamic studies and structural interpretation. *Protein Eng*, **10**, 541-549.

References

- [170] Fersht, A.R. (1985) *Enzyme structure and mechanism*, 2 ed. W.H. Freeman, New York.

